

Cardioversion

Cardioversion is the conversion of the heart rhythm from Atrial Fibrillation (or Atrial Flutter) to the normal rhythm, known as sinus rhythm. Electrical cardioversion is also known as Direct Current Cardioversion (DCCV).

Electrical Cardioversion

This may sound terrifying, but it is very simple in principle and it is a highly effective treatment in carefully chosen patients. The idea is to use an electric shock to activate the whole heart at once. This prevents the perpetuation of Atrial Fibrillation. After the shock the normal heart beat (sinus rhythm) will be able to emerge.

The cardioversion itself involves linking you to an ECG monitor which is connected to the cardioverter/defibrillator. An injection of short acting anaesthetic or powerful sedation is given. You will then be asleep and/or totally unconcerned about the procedure.

Risks Involved:

- Slow heart rhythm (bradycardia) – usually very transient and at most needing treatment with an intravenous medicine (Atropine) or a short period of pacing (electrical stimulation of the heart to initiate heart beats) for a short time.
- Fast heart rhythm (such as ventricular tachycardia) which may need a follow-up shock before you regain consciousness.
- Stroke, which is very unusual if you have been fully anticoagulated before the procedure.
- Skin burns or irritation from the electrodes (patches) - this is unusual with modern patch electrodes but can happen more frequently with older metal paddle electrodes.
- Early reversion of the normal rhythm back

to Atrial Fibrillation – this may require a further shock (when still under anaesthetic/sedation).

- General anaesthetic risks – rare in normal sized people with no other medical problems.

Electrode patches or plates are positioned on the back and front of the chest, or on the upper right and lower left of the chest. The cardioverter/defibrillator is charged and set to deliver a shock simultaneously with the next heart beat. Often the first shock is successful but sometimes several shocks at increasing energy levels or with different electrode patch positions is needed to convert the rhythm.

The normal rhythm is restored in about 90% of patients, but a small proportion immediately return to Atrial Fibrillation. Over the next few days, 10% - 20% lapse back into the arrhythmia but this can be reduced when necessary by prescription of an anti-arrhythmic drug.

After the procedure you will be awake within a minute or so and, although groggy for a while, you will quickly regain full control and will be ready to go home after a few hours. The ECG is monitored until you are fully recovered, a 12-lead ECG is recorded and you are then allowed to get up and move around. A friend or partner should come to hospital with you as you drive for 24 hours after the procedure and should be accompanied home. Someone should also stay with you on the night after the procedure in case they have a late complication.

Author: Professor A John Camm, EP
Endorsed by: Dr Matthew Fay, GP
Mrs Jayne Mudd, Arrhythmia Nurse Specialist
Published January 2009
Reviewed April 2010

For further information contact Atrial Fibrillation Association

Medical Advisory Committee: Dr Michael Davis,
Professor Ben Freedman, Dr Gerry Kaye